LIMIT

A limit's a dream in sight, We chase it with all our might! But just as we draw near, It grins, and disappears.

In mathematics, a limit is the value that a function or a sequence approaches as the input approaches some value. Limits are essential to calculus and mathematical analysis, and are used to define continuity, derivatives, and integrals.

3.1 DEFINITION OF LIMIT

Consider a function f(x) that is defined in a domain D which includes the point c. The function may or may not be defined at c. If, for all x that is close to c except for c, f(x) is arbitrarily close to a number L (as close to L as we like), then it is said that f approaches the limit L as xapproaches c and is written as:

$$\lim_{x \to c} f(x) = L$$


If the function can be evaluated at c, the limit L is simply f(c). But there can be situations where the function is not evaluable at c. E.g., the following function cannot be evaluated at x = 1.

$$\lim_{x \to 1} \frac{x^2 + x - 2}{x - 1}$$

the function is not evaluable at
$$c$$
. E.g., the following function of $\lim_{x\to 1}\frac{x^2+x-2}{x-1}$
But this function can be easily simplified to:
$$f(x)=\frac{(x-1)(x+2)}{x-1}=x+2$$

$$\to \lim_{x\to 1}f(x)=3.$$

FORMAL DEFINITION OF LIMIT 3.2



Let f(x) be a function that is defined on an interval that contains x = c, except possibly at c. Then, $\lim_{x\to c} f(x) = L$ if for every number $\epsilon > 0$, there is some number $\delta > 0$ such that, when $0 < |x - a| < \delta$, $|f(x) - L| < \epsilon$.

This means that for any number $\epsilon > 0$ that we pick, one can go to the graph and sketch two horizontal lines at $L + \epsilon$ and $L - \epsilon$. Then there must be another number $\delta > 0$ that can be determined to enable us to add in two vertical lines in the graph $a + \delta$ and $a - \delta$.

CHAPTER 3. LIMIT 13

3.3 Laws of Limit

Given L, M, c, k are real numbers such that $\lim_{x\to c} f(x) = L$ and $\lim_{x\to c} g(x) = M$. Then,

 $\lim_{x \to c} (f(x) + g(x)) = L + M$ Sum Rule $\lim_{x \to c} (f(x) - g(x)) = L - M$ Difference Rule

Constant Rule $\lim (kf(x)) = kL$ $\lim_{x \to \infty} (f(x)g(x)) = LM$ **Product Rule**

Ouotient Rule

 $\lim_{x \to c} \frac{f(x)}{g(x)} = \frac{L}{M}$ $\lim_{x \to c} [f(x)]^n = L^n \ (n > 0)$ Power Rule $\lim_{x \to \infty} \sqrt[n]{(f(x))} = \sqrt[n]{(L)} (n > 0)$ Root Rule

Examples:


$$\lim_{x \to 3} \sqrt{(2x^3 + 10)} = 8$$

$$\lim_{x \to 0} \frac{\sqrt{x^2 + 9} - 3}{x^2}$$

The above function is not evaluable at x = 0. The standard trick is to multiply both numerator and denominator by the conjugate radical expression.

$$\frac{\sqrt{x^2+9}-3}{x^2} = \frac{\sqrt{x^2+9}-3}{x^2} \cdot \frac{\sqrt{x^2+9}+3}{\sqrt{x^2+9}+3} = \frac{1}{\sqrt{x^2+9}+3} = \lim_{x \to 0} \frac{1}{\sqrt{x^2+9}+3} = \frac{1}{6}$$

3.4 AN IMPORTANT LIMIT



Consider the circle with a unit radius.

Area \triangle OAP < area sector OAP < area \triangle OAT

 $\frac{1}{2}sin \theta \le \pi 1^2 \left(\frac{\theta}{2\pi}\right) \le \frac{1}{2}tan \theta$ (θ is in radians)

$$1 \le \frac{\theta}{\sin \theta} \le \frac{1}{\cos \theta}$$

$$\rightarrow 1 \ge \frac{\sin \theta}{\theta} \ge \cos \theta$$

Hence,

$$\lim_{\theta \to 0} \frac{\sin \theta}{\theta} = 1 \qquad \text{where } \theta \text{ is in radians} \qquad (3.4.1)$$

Now consider the function $f(\theta) = \frac{1}{\sin \theta}$. Does it have a limit as $t \to \theta$ from either side? As θ approaches 0, its reciprocal, 1/x, grows without bound and the values of function cycle repeatedly from -1 to 1. There is no single number L that the function values stay increasingly close to as $\theta \to 0$. The function has neither a right-hand limit nor a lefthand limit at $\theta = 0$.

CHAPTER 3. LIMIT

3.5 One Sided Limits

$$\lim_{\substack{x \to 0^{+} \\ \lim_{x \to 0^{-}} f(x) = -1}} f(x) = 1$$

3.6 Continuous Function

Function is right-continous at c (continuous from right) if $\lim_{x\to c^+} f(x) = f(c)$ Function is left-continous at c (continuous from left) if $\lim_{x\to c^-} f(x) = f(c)$ A function is continous at c if $\lim_{x\to c} f(x) = f(c)$


If a function is discontinuous at one or more points of its domain, it is called a discontinuous function.

3.7 Infinite Limits

$$\lim_{x\to 0^+} \frac{1}{x} = \infty$$
, $\lim_{x\to 0^-} \frac{1}{x} = -\infty$ Note that this does not mean that the limit exists as there is no real number such as ∞ . It is simply a concise way of saying that the limit does not exist.

3.8 Rolle's Theorem


If f is a continuous function on a closed interval [a,b] and If f(a)=f(b), then there is at least one point c in (a,b) where $f^{'}(c)=0$.



3.9 Mean Value Theorem

There is at least one number c in the interval (a,b) such that:

$$f^{'}(c) = \frac{f(b) - f(a)}{b - a}$$



CHAPTER 3. LIMIT 15

3.10 SymPy Code

Determine the limits of:

$$\lim_{x \to 1} \frac{x^2 + x - 2}{(x - 1)}$$

$$\lim_{x \to 0} \frac{\sqrt{x^2 + 9} - 3}{x^2}$$

$$\lim_{x \to 0} sin(x)$$

```
import sympy as sp
  import numpy as np
  import matplotlib.pyplot as plt
  from IPython.display import display, Math
  from sympy import sin, cos, tan, trigsimp, expand_trig
  from sympy import oo
  from sympy import limit
  x = sp.symbols('x')
  y = (x**2 + x - 2) / (x - 1)
lim = limit(y, x, 1)
  display(lim)
14 y = ((x**2 + 9)**0.5 - 3) / x**2
  lim = limit(y, x, 0)
  display(lim)
18 y = \sin(x)/x
  \lim = \lim(y, x, 0)
  display(lim)
```

```
\frac{1}{6}
\cos(x)
```

If you are copying code from pdf into your editor or python ternminal, please replace the following in your editor. Regex $\space^*\s$