Integration

Integration sounds lofty and grand, Yet it's just adding, hand over hand. Cloaked in fine symbols, solemn, sublime -Still it's just summing, one step at a time!

Integration, in mathematics, is the technique of finding a function g(x) the derivative of which is equal to a given function f(x). This is indicated by the integral sign \int as in $\int f(x)dx$ and is called the indefinite integral of the function. The symbol dx represents an infinitesimal displacement along x. Hence, $\int f(x)dx$ is the summation of the product of f(x) and dx. The definite integral, written as $\int_a^b f(x)dx$ where a and b are called the limits of integration, is equal to g(b) - g(a), where $\frac{d}{dx}g(x) = f(x)$.

5.1 Integral

Given a function f(x), an anti-derivative of f(x) is any function g(x) such that g'(x) = f(x). The most general anti-derivative is called the indefinite integral.

$$\int f(x)dx = g(x) + c$$
 where c is a constant of integration

Note the following inequalities.

$$\int f(x)g(x)dx \neq \int f(x)dx \int g(x)dx$$

$$\int \frac{f(x)}{g(x)} dx \neq \frac{\int f(x) dx}{\int g(x) dx}$$

5.2 Common Integrals

$$\int x^n dx = \frac{x^{(n+1)}}{(n+1)} + c \tag{5.2.1}$$

$$\int e^x dx = e^x + c \tag{5.2.2}$$

$$\int a^x dx = \frac{a^x}{\ln a} + c \tag{5.2.3}$$

$$\int \frac{1}{x} dx = \ln|x| + c \tag{5.2.4}$$

$$\int \cos(x)dx = \sin(x) + c \tag{5.2.5}$$

CHAPTER 5. INTEGRATION 24

5.3 Substitution Technique

$$\int 18x^{2} \sqrt[4]{(6x^{3} + 5)} dx$$
Let $u = 6x^{3} + 5$

$$\to du = 18x^{2} dx$$

$$\to \int \sqrt[4]{u} du = \frac{u^{\left(\frac{1}{4} + 1\right)}}{\frac{1}{4} + 1} = \frac{4}{5} u^{\frac{5}{4}} = \frac{4}{5} \left(6x^{3} + 5\right)^{\frac{5}{4}}$$

5.4 Integration by Parts

$$[f(x)g(x)]' = f(x)g'(x) + f'(x)g(x)$$

$$f(x)g'(x) = [f(x)g(x)]' - f'(x)g(x)$$

$$\int f(x)g'(x) dx = \int [f(x)g(x)]' dx - \int f'(x)g(x) dx$$


$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx$$

$$\int u dv = uv - \int v du$$
(5.4.1)

Hence, integral of two functions = first function \times integral of second function — integral of (differntiation of the first function \times integral of the second function).

5.5 Definite Integral

A definite integral is a the area under its curve.



$$\left| \int_a^b f(x)dx = \lim_{n \to \infty} \sum_{i=1}^n f(x_i^*) \Delta x = g(x) \right|_a^b = g(b) - g(a)$$

where $f(x_i^*)$ is the value at the middle of the strip Δx .

$$f_{avg} = \frac{1}{b-a} \int_{a}^{b} f(x) dx$$

$$\int_a^b f(x) dx = f(c)(b-a) \text{ where c is in [a,b]}$$

CHAPTER 5. INTEGRATION 25

5.6 Some Integration Strategies

- (1) Simplify the integrand. E.g., $cos^2(\theta) = \frac{1}{2}(1 + cos(2\theta))$
- Check if simple substitution will work
- 3 If integrand is a rational expression, partial functions may work
- 4 If integrand is polynomial x, trig, exp, ln function, integration by parts may work
- 5 If integrand involves $\sqrt{b^2x^2 + a^2}$, trigonometric substitution may work
- 6 If integrand has a quadratic in it, completing the square may work.

5.7 SymPy

Integrate the following functions:

$$\int (x^2 + x + 1) dx$$

$$\int e^{-x^2} \operatorname{erf}(x) dx \text{ where } \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$$

$$\int y = e^x dx$$

```
import sympy as sp
  import numpy as np
  import matplotlib.pyplot as plt
  from IPython.display import display, Math
  from sympy import sqrt, diff, integrate, oo
  from sympy import sin, cos, tan, ln, exp, erf, trigsimp, expand_trig, simplify
  from sympy import sinh, cosh
  x = sp.symbols('x')
  y = x**2 + x + 1
  int_expr = integrate(y,x) # integrate y wrt x
  print(sp.latex(int_expr))
14 display(int_expr)
16 y = \exp(-x**2)*\operatorname{erf}(x)
int_expr = integrate(y,x)
                                          # integrate y wrt x
18 print(sp.latex(int_expr))
19 display(int_expr)
y = \exp(-x)
int_expr = integrate(y, (x, 0, oo)) # definite integral, limits 0 & infinity
print(sp.latex(int_expr))
24 display(int_expr)
```

$$\frac{x^3}{3} + \frac{x^2}{2} + x$$

$$\frac{\sqrt{\pi}\operatorname{erf}^2(x)}{4}$$